

REVIEW ARTICLE

Overweight and type II diabetes among the elderly in Nigerians Belt

Dr. Slim A, afrin^{1*} | Dr. M Ahmad²

¹Department of Chemical Pathology, College of Health Science, University of Jos, Jos, Nigeria

²Department of Chemical Pathology, College of Medicine, University of Jos, Jos, Nigeria

Abstract

Background: The increasing rate of obesity and type-2 diabetes (T₂DM) are major impending threats to the health of African Population but the magnitude of the two in middle age Nigerians is not known. We assessed the burden of obesity and T₂DM among elderly middle belt Nigerians.

Methods: A cross sectional prospective study was conducted among the population living in Lafia town, the capital of Nasarawa State in Middle Belt Nigeria.

After biodata and clinical examination, including arthrometric measurements of weight and height was conducted. Standard BMI was calculated. A casual glucose and fasting blood glucose was determined using serum enzymatic glucose oxidase method. The diagnostic criteria were taken as fasting glucose at $\geq 7.0\text{mmol/L}$ and casual glucose at $\geq 11.1\text{mmol/L}$.

Results: In Lafia town the prevalence of Class I obesity was 62.6%. There is a perfect correlation of 1.000 between BMI and glucose, meaning that as the BMI of a person increases, blood glucose concentration always increases and vice versa. The correlation between age and glucose concentration (0.0447) reflects a very weak positive relationship, indicating that BMI may differ at different ages. i.e. as the age of a patient increases, there is a slight tendency that the BMI may also increase.

Conclusion: This study showed high risk of T₂DM and obesity among middle belt Nigerians living in Lafia town due to sedentary lifestyle and lack of exercise in addition to indiscriminate diet. There is need for regular screening for diabetes and hypertension. Use of portable electronic BP machine is advocated. There should be lifestyle modification like regular exercise, control of excessive alcohol intake and dietary advice. Keywords: T2DM, BMI, obesity, BP

1 | INTRODUCTION:

Overweight and especially obesity, particularly at young ages, substantially increase risk of diagnosed diabetes while the impact on diabetic risk life expectancy duration diminishes with age.

Type 2 diabetes mellitus (T₂DM) comprises about 80-94% of all diabetes cases and its prevalence has been steadily increasing. Obesity, classified as body mass index (BMI) $\geq 30\text{kg/m}^2$, is a known predictor of T₂DM and has become a major public health problem in the US, affecting over one-third (35.7%) of the population. The rise in prevalence of obesity among all age diet, and urbanization-explain the rising of diagnosed type-2 DM (1–16)

There is evidence that both genetic and environmental factors contribute in the pathogenesis of type-2 DM.

Numerous studies have shown that obesity (central obesity in particular), physical inactivity, high-fat diet, and diet rich in saturated fatty acids increase the risk of diabetes. Based on epidemiological studies, low intakes of dietary fiber, low-glycemic carbohydrates, and whole grain cereals have been shown to increase the risk of type-2 DM

It is clear that increased body weight is a risk factor for type-II DM, the relationship between body weight and Type is more properly attribute to the quantity and distribution of body fat (adiposopathy). Increase adipose tissue distribution leads to peripheral insulin resistance leading to T₂DM. That is why decreasing weight proportionally lower blood glucose levels.

The clinical value of measuring BMI from a diabetes perspective lies the whether this measure can identify individuals who may have undiagnosed diabetes or may be at increased future risk for diabetes. In addition measuring BMI also is important for managing diabetes for the purpose of weight control^{4–6}.

BMI cut off have been established to identify overweight (BMI $\geq 25\text{kg/m}^2$) or obese (BMI $> 30\text{kg/m}^2$ individual).

The prevalence of excess body weight has increased dramatically in the United States in the past 50 years.

In 2010, 69.2% of the US adult population had a body mass index (BMI) above normal levels. Among adults older than age 60, the prevalence of excess weight was even higher. Adult obesity is predicted to rise by 33% in the next 2 decades, with a prevalence of severe obesity rising by 130%.

A study in Ghana showed a high risk of obesity and T₂DM among sub-Sahara Africa living in Europe⁹.

In Nigeria, puppet et al found over-weight to be 22.6% prevalence in Jos and appear to the increasing. This presence study is aimed at assessing the present effect of BMI1 on diabetes in Jos¹⁰.

This study is aimed at exploring the threshold and onset of Diabetes mellitus at various age group.

2 | MATERIALS AND METHOD

Sampling Data Collection and Analysis

This is a cross sectional prospective study covering Lafia metropolis and all the neighboring villages. Seven Hundred (700) elderly participants mostly retired civil servants in Lafia with age range of 40-90 years.

Both males and females were recruited into study. Physical examination was carried out by the authors. The weights of the subjects were measured to the nearest kilogram with a Hanson type bathroom weighing scale. The height was measured to the nearest centimeter. The body mass index (BMI) was calculated and recorded. Classification of BMI was done according to the recommendations of the WHO expert committee for the classification of overweighting¹⁰.

Supplementary information The online version of this article contains supplementary material, which is available to autho-rized users.

Corresponding Author: Dr. Slim A, afrin
Department of Chemical Pathology, College of Health Science, University of Jos, Jos, Nigeria

Venous blood (2.5ml) was collected into fluoride oxalate and transported to the chemical pathology laboratory in Shalom Hospital. The WHO diagnostic criteria of casual blood glucose level $\geq 11.1\text{mmol/L}$ and fasting glucose value $\geq 7.0\text{mmol/L}$.

Statistical Analysis

The data obtained were coded and entered into Stata Software for analysis. The data are presented as mean \pm S.D. comparison was done by student's t-test for continuous variables.

Ethical Consideration

The study was conducted with adherence to ethical standards; informed consent was used in the recruitment of participants. Approval for this study was obtained from Ethics Committee of JUTH; confidentially was maintained in accordance with standard medical practice.

3 | RESULTS

The study shows result of overweight and diabetes mellitus in this elderly population.

The glucose (mmol/L) levels and the weight (kg) and height (m), ages (years) Blood pressure (mmHg) and gender of all the participants reveal this results below.

TABLE 1: Correlation OF BMI, Glucose with age and gender

	BMI	Diabetes	
BMI	0.1183	1.000	
Glucose	-0.0236		
Age	0.1284	0.0447	-0.0593
Gender	-0.0678	0.0149	-0.0274

From table one above, there is a perfect correlation of 1.000 between BMI and glucose, this means that as the BMI of a person increases, a glucose concentration always increases also.

The correlation between age and glucose concentration (0.0447) reflects a very weak positive relation-

ship, indicating that BMI may differ at different ages. i.e. as the age of a patient increases, there is a slight tendency that the BMI may also increase.

TABLE 2: BMI Class and population

Weight (kg)	BMI Class	Population	% Population
Underweight	<18	37	4.03
Normal weight	18-24	224	24.43
Overweight	25-29	574	62.60
Class I Obesity	30-34	50	5.4
Class II Obesity	35-39	28	2.7
Class III Obesity	>40	3	0.53

From the table above overweight have the highest percentage population of 62.6 Class I (2.73%) with least in Class II (0.3%).

TABLE 3: Glucose levels and population

Glucose (mmol/L)	Population	% Population
1-5	100	10.91
6-10	230	25.08
11-15	78	8.51
16-20	6	0.65

In table 3 the glucose concentration in mmol/L was classified into new diabetic range of 1-5mmols and the prediabetes and diabetic range i.e. 6-10, 11-15.

The coefficient of variation for the glucose estimations was 6.4% at 5.6mmol/L and 2.1% at 18.3mmol/L.

The highest population falls into the range of 6-10mmol/L and the diabetic range is 11-15mmol giving us 8.51% population as diabetics.

Diagnosed diabetes fallen as casual glucose of 11.1mmol/L

Histograms

4 | DISCUSSION

This study showed a perfect correlation between BMI and diabetes in this population. It shows BMI (30kg/m^2) to be 7.1% prevalence. The highest percentage (62.6%) population falls in the class of overweight ($25\text{-}29\text{kg/m}^2$).

Recently, the United States centre for disease control showed a prevalence of obesity in Asian-America was only 10.8% compared with 34.9% in all US adults in 2011-2012¹². The prevalence of excess body weight has increased dramatically in the United States in the past 50 years¹²⁻¹⁴.

This value of 7.1% is a warning sign from this study. BMI cutoff have been established to identify overweight ($\text{BMI} \geq 25\text{kg/m}^2$) or obese ($\text{BMI} \geq 30\text{kg/m}^2$) individuals.

Adjusting for a number of characteristics associated with the risk of T₂DM, we found that, compared with normal BMI, overweight and obesity was statistically significantly associated with the risk of being diagnosed with T₂DM among individuals without any other prior evidence of T₂DM.

We further found that the risk of a T₂DM diagnosis was increasing larger for individuals in higher BMI categories than for individuals in lower BMI categories¹⁵⁻¹⁷.

These results agree with other studies that have examined the association between BMI and risk of T₂DM using nationally representative samples. For example, using data from the 2001 Behavioral Risk Surveillance System.

Furthermore, found statistically significant and increasingly larger ORs and T₂DM among overweight adults (1.59, 95% CI: 1.46-1.73) adults with BMI between 30 and 39.9kg/m^2 (3.44, 95% CI: 3.17-3.74), adults with $\text{BMI} \geq 40\text{kg/m}^2$ (7.37, 95% CI: 6.39-8.50) relative to adults with normal BMI (20)¹⁸⁻²⁰.

Moreover, BMI values were clinically measured in the current study, compared with BMI calculated from self-reported height in those earlier studies. Self reported weight and height considerably underestimate the individuals measured BMI^{31,32} and many thus have weakened the association between obesity and risk of T₂DM and/or biased the estimated results.

This may explain the lower ORs associated with BMI levels in the Mokdad, *et al* study compared to the current study²¹⁻²³.

The risk of developing T₂DM for individuals who were overweight or obese was about 1.5-5 times higher than for individuals with normal BMI, as estimated in our study. This demonstrates the importance of continuous weight management, which not only can reduce the disease burden of obesity but also may prevent further progression to T₂DM weight management is particularly important for people with severe obesity who never disproportionately at higher risk²⁴⁻²⁵.

The influence of time and modernization may explain the difference between these two studies. Advancing age was another identified independent risk factor for diabetes, which was more prevalent in subjects aged 50years and above. The TallyHo mouse model, shows diabetic characteristics similar to those of humans with diabetes. Johnson and McLarty *et al* found that the peak incidence of diabetes in Nigeria and Tanzania, respectively was after 45-50years of age. It is well known that the prevalence of diabetes increases with age¹⁹. There was increased body weight, increased blood glucose levels, abnormal insulin. These features found to increase with age in the TallyHo mice, just as they were found to increase with age in humans with diabetes. This increase in age as a risk factor was also found to agree with this work. In Nigeria, the risk of diabetes increases 3-4 folds after the age of 44years¹¹. The worsening of insulin resistance with age and increasing longevity of diabetic patients due to improved care, all contribute to the rising prevalence of type-2 diabetes with age¹⁴⁻¹⁹.

Physical inactivity is a well known risk factor for type-2 diabetes. The risk of diabetes is reduced by 50% among men who take moderately vigorous exercise²⁶. This suggests that not all diabetes is the same or that other factors markedly influence its impact on health.

The cardiovascular sequelae of diabetes may differ, depending on whether the diabetes is hyperinsulinemic, early or late in onset caused by some known cause, or treated or untreated.

Some studies of cigarette smoking among diabetics have been punished^{20–22}. The finding of reduced cigarette consumption among diabetic males appears to be unique.

In general, it does not appear that diabetics cope less well with risk factors than do non-diabetics. Although the present analysis is based on a limited number of diabetics, the evidence of consistently against interaction between diabetes and cardiovascular risk factors.

REFERENCES

1. Vessby B, Uusitupa M, Hermansen K, Riccardi G, Rivellese AA, Tapsell LC, et al. Substituting dietary saturated for monounsaturated fat impairs insulin sensitivity in healthy men and women: The KANWU study. *Diabetologia*. 2001;44(3):312–319. Available from: <https://dx.doi.org/10.1007/s001250051620>. doi:10.1007/s001250051620.
2. Tesfaye F, Minh PV, Byass Y, Berhane R, Bonita NG, Nawi et al. association between BMI and blood pressure across three populations in Africa and Asia. *Am J Heart Assoc*. 2017;(3):501–504.
3. Groop LC. Insulin resistance. The trigger of type-2 diabetes. obesity and metabolism. 1999;1:1–7.
4. Gordon T, Garcia-Palmieri MR, Kagan A, Kannel WB, Schiffman J. Differences in Coronary Heart Disease in Framingham, Honolulu and Puerto Rico. *Journal of Chronic Diseases*. 1974;27(7-8):329–344. Available from: [https://dx.doi.org/10.1016/0021-9681\(74\)90013-7](https://dx.doi.org/10.1016/0021-9681(74)90013-7). doi:10.1016/0021-9681(74)90013-7.
5. Kim JH, Šaunak Sen, Avery CS, Simpson E, Chandler P, Nishina PM, et al. Genetic Analysis of a New Mouse Model for Non-Insulin-Dependent Diabetes. *Genomics*. 2001;74(3):273–286. Available from: <https://dx.doi.org/10.1006/geno.2001.6569>. doi:10.1006/geno.2001.6569.
6. Hu FB, Liu S, van Dam RM. Diet and risk of Type II diabetes: the role of types of fat and carbohydrate. *Diabetologia*. 2001;44(7):805–817. Available from: <https://dx.doi.org/10.1007/s001250100547>. doi:10.1007/s001250100547.
7. Zimmet P. Type 2 (non-insulin-dependent) diabetes ? An epidemiological overview. *Diabetologia*. 1982;22(6):399–411. Available from: <https://dx.doi.org/10.1007/bf00282581>. doi:10.1007/bf00282581.
8. Rimm EB, Chan J, Stampfer MJ, Colditz GA, Willett WC. Prospective study of cigarette smoking, alcohol use, and the risk of diabetes in men. *BMJ*. 1995;310(6979):555–559. Available from: <https://dx.doi.org/10.1136/bmj.310.6979.555>. doi:10.1136/bmj.310.6979.555.
9. WHO Expert Committee on diabetes mellitus Second Report WHO Technical Report Series. 1980;646.
10. Stevens J, Cai J, Pamuk ER, Williamson DF, Thun MJ, Wood JL. The Effect of Age on the Association between Body-Mass Index and Mortality. *New England Journal of Medicine*. 1998;338(1):1–7. Available from: <https://dx.doi.org/10.1056/nejm199801013380101>. doi:10.1056/nejm199801013380101.
11. Singh RB, Rastogi SS, Rastogi V, Niaz MA, Madhu SV, Chen M, et al. Blood pressure trends, plasma insulin levels and risk factors in rural and urban elderly populations of North India. *Coronary Artery Disease*. 1997;8(7):463–468. Available from: <https://dx.doi.org/10.1097/00019501-199707000-00009>. doi:10.1097/00019501-199707000-00009.
12. Shurtleff D. DHEW Publication No;.
13. Ganz ML, Wintfeld N, Li Q, Alas V, Langer J, Hammer M. The association of body mass index with the risk of type 2 diabetes: a case-control study nested in an electronic health records system in the United States. *Diabetology & Metabolic Syndrome*. 2014;6(1):50–50. Available from: <https://dx.doi.org/10.1186/1758-5996-6-50>. doi:10.1186/1758-5996-6-50.

14. Gray N, Picone G, Sloan F. Arseniy Yashkin. Relation between BMI and Diabetes Mellitus and its complications among US older adults. South Med J. 2015;108(1):29–36.
15. West KM. Diabetes in American Indians and Other Native Populations of the New World. Diabetes. 1974;23(10):841–855. Available from: <https://dx.doi.org/10.2337/diab.23.10.841>. doi: 10.2337/diab.23.10.841.
16. Mclarty DG, Swai AB, Kitange HM, Masuki G, Mitnangi BL, Kilama PM, et al. prevalence of diabetes and impaired glucose tolerance in rural Tanzania. 1989;1:871–875.

How to cite this article: Afrin D.S., Ahmad D.M. Overweight and type II diabetes among the elderly in Nigerians Belt. Journal of Medical Reviews. 2020;6–10.
